

Transfer of Leachable Substances from Cross-Linked Peroxide Polyethylene (PEX-a) into Hemodialysis Water

Faissal Tarrass¹ , Meryem Benjelloun¹ 

¹ Center of Hemodialysis 2 Mars, Casablanca, Morocco

Contributorship Statement:

- CO, AM: Were involved in drafting and editing the manuscript.
- CO: Revised the manuscript prior to submission.

Keywords: Dialysis Solutions; Polyethylene Glycols; Renal Dialysis; Water

Cross-linked polyethylene, commonly abbreviated PEX or XLPE, is a thermoplastic polymer, which is obtained through the polymerization of the ethylene.¹ Cross linking of polyethylene into PEX pipes results in improved properties, such as thermal, chemical and abrasion resistance, as well as a slower crack growth, accompanied with a greater toughness.¹ PEX piping is commonly used in home plumbing systems for heating, cooling, and water distribution systems.¹ Recently, polyethylene cross linked using peroxide (PEX-a) has become a viable alternative

to polyvinyl chloride (PVC), or chlorinated polyvinyl chloride (C-PVC) for use as distribution systems for ultra-pure hemodialysis water.²

Research has shown that PEX-tubing can leach organic carbon and sorbing contaminants into water, especially when heated,³ exposed to high PH levels,⁴ chlorine-based disinfectants⁴ or ultra-violet (UV) irradiation.⁵ More than 158 compounds including antioxidants and their degradation products, compounds of known origins, and compounds with unknown origins were detected and identified in water which was in contact with PEX-a tubing.⁶ Some examples include

Table 1

Chemical substances potentially leaching from PEX-a

Chemical group	CAS Number	Substance	Reference
Acetophenone derivatives	1378888-43-7	1-(3-Ethyl-4-(hydroxymethyl) phenyl)ethanone	7
	30773-71-8	1,1'-(Phenylene)bisethanone	7
Alcohols	110-03-2	2,5-dimethyl-2,5-hexanediol	7,8
	104-76-7	2-Ethyl-1-hexanol	8,9
	67-56-1	Methanol	7
	111-87-5	n-Octanol	9
	75-65-0	tert-Butyl alcohol	7,11
Aldehydes	107-86-8	3-Methyl-2-butenal	7
	112-31-2	Decanal	8,9
	124-19-6	Nonanal	6,11
Alkanes	1560-93-6	2-Methyl pentadecane	12
	2882-96-4	3-Methyl pentadecane	12
	629-78-7	n-Heptadecane	12
	544-76-3	n-Hexadecane	12
	629-62-9	n-Pentadecane	12
Alkyl phenol	128-37-0	Butylatedhydroxytoluene	7,11
	20170-32-5	3-(3,5-Di-tert-butyl-4- hydroxyphenyl)propanoic acid	9
	6386-38-5	Methyl 3-(3,5-di-tert-butyl-4-hydroxyphenyl)propionate	7,9,12
	-	Cyclohexa-1,4-diene-1,5- bis(tert-butyl)ethylidene	13
	1620-98-0	3,5-Di-tert-butyl-4- hydroxybenzaldehyde	9,13
	14035-33-7	3,5-Di-tert-butyl-4- hydroxyacetophenone	13
	96-76-4	2,4-Di-tert-butylphenol	10
	128-39-2	2,6-Di-tert-butylphenol	10
	121-00-6	2-tert-Butyl-4-methoxyphenol	14
	122-94-1	4-Butoxyphenol	11
	-	Phenolics	7

© Author(s) (or their employer(s)) and Portuguese Journal of Nephrology & Hypertension 2023. Re-use permitted under CC BY 4.0. (<https://creativecommons.org/licenses/by/4.0/>).

Chemical group	CAS Number	Substance	Reference
Amides	–	Diazadiketo-cyclo-tetradecane	7
Aniline derivatives	101-67-7	4,4'-Dioctyldiphenylamine	12
Aromatic hydrocarbons	1014-60-4	1,3-Di-tert-butylbenzene	12
	1012-72-2	1,4-Di-tert-butylbenzene	7
	71-43-2	Benzene	7,11
	110-82-7	Cyclohexane	12
	95-47-6	o-Xylene	12
	106-44-4	p-Cresol	12
	106-42-3	p-xylene	12
	108-88-3	Toluene	9,11,12
	1330-20-7	Xylene(s)	8,11
Biolefinicsteroid	1224-94-8	Androsta-5,16-dien-3beta-ol	12
Bisphenol compounds	80-05-7	Bisphenol A	7
Epoxides	106-92-3	Oxirane, ((2-propenyloxy)methyl)-	7
Esters	103-23-1	Bis(2-ethylhexyl) hexanedioate	12
	109-21-7	Butyl butyrate	7
	540-88-5	tert-Butylacetate	9
	77-93-0	Triethyl citrate	12
	629-82-3	1-Oxtoxyoctane	9
	126-84-1	2,2-Diethoxypropane	8
Ethers	637-92-3	Ethyltert-butylether	8,12
	33021-02-2	Isopropyltert-butylether	11
	1634-04-4	Methyltert-butylether	11
Fatty acids	334-48-5	Decanoicacid	9
	143-07-7	Dodecanoicacid	9
	124-07-2	Octanoicacid	9
Halogenated hydrocarbons	598-99-2	Methyltrichloroacetate	9
	630-20-6	1,1,1,2-Tetrachloroethane	9
	79-01-6	Trichloroethylene	4
Heterocycles	15045-43-9	2,2,5,5-Tetramethyloxolane	6
	95-16-9	Benzothiazole	4
	110-86-1	Pyridine	9
	109-99-9	Tetrahydrofuran	4
Hydrocarbons	74663-85-7	Nonylcyclopropane	4
Inorganic	75-15-0	Carbon disulfide	4
Ketones	110-12-3	5-Methyl-2-hexanone	8
	108-94-1	Cyclohexanone	4
	120-92-3	Cyclopentanone	4
	–	Dicyclopentylone	4
	141-79-7	Mesityl oxide	8
Peroxides	110-05-4	Di-tert-butylperoxide	7
Phthalates	–	Butyl-2-methoxyethylphthalate	9
Quinones/quinone derivatives	106-51-4	1,4-Benzoquinone	4
	719-22-2	2,6-Di-t-butyl-p-benzoquinone	6-8
	82304-66-3	7,9-Di-tert-butyl-1-oxaspiro(4,5)deca-6,9-diene2,8-dione	4,9
Vinyl compounds	763-32-6	3-Methylbut-3-en-1-o	6
	115-11-7	Isobutylene	4,7
	115-18-4	Methylbutenol	4

CAS: Chemical Abstracts Service

phenols,⁷⁻¹⁴ aldehydes,^{7,8,11} esters,^{8,9,12} alcohols,^{7-9,12} ethers,^{8,11,12} hydroperoxides,¹⁰ ketones,^{7,11} phthalates,¹² and organic carbon, which could potentially act as substrate for the growth of microorganisms on the pipe surface.⁴ In a thesis work, Durand has found an increase in organic carbon for water exposed to PEX-a by 0.4 mg/L in the presence of chlorine based disinfectants.⁴ Also, in another work, Connell *et al*, has found that the leached organic carbon from PEX-a was greater

than 0.15 mg/L after 7 days at 23°C,¹² exceeding the threshold value (0.1 mg/L > 15°C) which significantly increased the probability of coliform occurrence.¹⁵ Table 1 report the common chemical substances leaching from PEX-a.

In hemodialysis, PEX-a tubing used as water distribution systems are raising concern about the potential for leaching chemicals into

the dialysis water, especially when heated during the thermal disinfection or exposed to chlorine-based disinfectants.¹⁶ Patients on hemodialysis are exposed to hundreds of liters of dialysis water each week, however there are currently no guidelines regarding acceptable levels of these leachable substances in the dialysis water. The health effects of the leaching chemicals that have been evaluated toxicologically are significant, ranging from liver and kidney effects to adverse health outcomes on the reproductive, developmental, immune and nervous systems, endocrine disruption, and/or carcinogenicity.¹⁷ To date, no studies had been conducted to ascertain the effects of PEX-a leaching on dialysis water quality. We propose that particular attention be paid to research on this field in order to ensure patient safety.

References

- Jinca T, Sabu T, Ahmad Z. Crosslinkable Polyethylene: Manufacture, Properties, Recycling, and Applications. Singapore: Springer Nature; 2021.
- Bolasco P. The production of on-line dialysis water for extracorporeal dialysis: proposals for an increased safety upgrade, a viewpoint. *J Nephrol.* 2020; 33: 405–15. doi: 10.1007/s40620-019-00667-2.
- Holder SL, Hedenqvist MS, Nilsson F. Understanding and modelling the diffusion process of low molecular weight substances in polyethylene pipes. *Water Res.* 2019; 157: 301-9.
- Durand M. Disinfectants and Plumbing Materials: Effects on Sensory and Chemical Characteristics of Drinking Water. [Thesis, Department of Environmental Engineering, Faculty of the Virginia Polytechnic Institute and State University]. Blacksburg: FVPISU; 2005.
- Anayo JM, Okpashi VE, Onwurah I. Impact of ultra violet radiation on polyethylene packaged water exposed at varying conditions: are we drinking micro-plastics? *Am J Biochem Biotech.* 2018; 14: 20-2
- Whelton AJ, Nguyen T. Contaminant migration from polymeric pipes used in buried potable water distribution systems: A review. *Crit Rev Environ Sci Technol.* 2013; 43: 679–751.
- Tomboulian P, Schweitzer L, Mullin K, Wilson J, Khiari D. Materials used in drinking water distribution systems: contribution to taste-and-odor. *Water Sci Technol.* 2004; 49: 219–26.
- Lin T, Watson S, Dietrich MA, Suffet IH. Taste and Odour in Source and Drinking Water: Causes, Controls, and Consequences. London: IWA Publishing; 2019.
- Loschner D, Rapp T, Schlosser FU, Schuster R, Stottmeister E, Zander S. Experience with the application of the draft European Standard prEN 15768 to the identification of leachable organic substances from materials in contact with drinking water by GC-MS. *Anal Methods.* 2011; 3: 2547–56.
- Denberg M, Mosbæk H, Hassager O, Erik A. Determination of the concentration profile and homogeneity of antioxidants and degradation products in a cross-linked polyethylene type A (PEXa) pipe. *Polymer Test.* 2009; 28: 378-85
- Skjevraak I, Due A, Gjerstad KO, Herikstad H. Volatile organic components migrating from plastic pipes (HDPE, PEX and PVC) into drinking water. *Water Res.* 2003; 37: 1912–20.
- Connell M, Stenson A, Weinrich L, LeChevallier M, Boyd SL, Ghosal RR, et al. PEX and PP water pipes: assimilable carbon, chemicals, and odors. *J Am Water Works Assoc.* 2016; 108: E192–E204.
- Brocca D, Arvin E, Mosbaek H. Identification of organic compounds migrating from polyethylene pipelines into drinking water. *Water Res.* 2002; 36: 3675–80.
- Lutzhoft HH, Waul CK, Andersen HR, Seredynska-Sobecka B, Mosbaek H, Christensen N, et al. HS-SPME-GC-MS analysis of antioxidant degradation products migrating to drinking water from PE materials and PEX pipes. *Int J Environ Anal Chem.* 2013; 93: 593–612.
- Volk C, LeChevallier M. Assessing biodegradable organic matter. *Am Water Works Assoc J.* 2000; 92: 64-76.
- Mime MI, Bellebia S, Bengharez Z, Louni M, Benrachedi K. Assessment and monitoring the quality of treated water and dialysate in a public hemodialysis center to prevent potential health risks. *Alger J Environ Sci Technol.* 2019; 5: 912-9.
- Barton C. Trichloroethylene. In: *Encyclopedia of Toxicology.* London: Elsevier; 2014. p. 827-30.

■ Ethical Disclosures

Conflicts of Interest: The authors have no conflicts of interest to declare.

Financial Support: This work has not received any contribution grant or scholarship.

Provenance and Peer Review: Not commissioned; externally peer reviewed.

Consent for Publication: Not applicable.

Corresponding Author:

Faissal Tarrass 
 Center of Hemodialysis 2 Mars
 466 Boulevard 2 Mars, Haddaouia 20460 Casablanca, Morocco
 Email: ftarrass@hotmail.com